arXiv:1510.00497v1 [math.DG] 2 Oct 2015

Global Poisson Structures on S^4

Takayuki MORIYAMA and Takashi NITTA

Abstract. A Poisson structure is a bivector whose Schouten bracket vanishes. We study a global Poisson structure on S^4 associated with a holomorphic Poisson structure on $\mathbb{CP}^3$. We provide a real structure and show that the space of the Poisson structures on S^4 is the real form of the space of holomorphic Poisson structures on $\mathbb{CP}^3$. We also obtain many examples of global Poisson structures on S^4.

1 Introduction

A global antisymmetric 2-vector is called a bivector. The Schouten bracket of two bivectors is defined by the convolution of the covariant derivative of the one and the another, further antisymmetrization of it. The bivector is called a Poisson structure if the Schouten bracket of the bivector with itself vanishes. Since the Schouten bracket is valued in the space of antisymmetric 3-vectors, any bivector is always a Poisson structure on 2-dimensional manifolds. In general, the equation for vanishing the Schouten bracket is written as a system of partial differential equations. We can find that it is very complicated to solve the equation even on 3-dimensional manifolds. Moreover, it is much harder to solve the equation globally on the manifold. Hence, on 3-dimensional manifolds, the existence of the Poisson structure is non-trivial and the construction of it is difficult.

In the holomorphic category, we consider holomorphic bivectors and holomorphic Schouten brackets (c.f. [3]). When the Schouten bracket vanishes, the holomorphic bivector is called a holomorphic Poisson structure. Since the Schouten bracket is also a holomorphic antisymmetric 3-vector, any holomorphic bivector is always a holomorphic Poisson structure on complex surfaces. Recently, Goto and Hitchin studied deformations of the holomorphic Poisson structures ([2], [4]).

In this paper, firstly we concentrate global holomorphic Poisson structures on the 3-dimensional complex projective space $\mathbb{CP}^3$. We show that the space of holomorphic Poisson structures on $\mathbb{CP}^3$ is an algebraic variety in Section 3. We also obtain some examples of holomorphic Poisson structures on $\mathbb{CP}^3$. It is known that holomorphic foliations induces holomorphic Poisson structures on $\mathbb{CP}^3$. The space of holomorphic foliations of degree 2 is completely classified and has the six irreducible components

2010 Mathematics Subject Classification. Primary 53D17.
Key Words and Phrases. Poisson structure, holomorphic Poisson structure, Schouten bracket.

[1]. Hence, we can provide a lot of examples of holomorphic Poisson structures on $\mathbb{CP}^3$.

Secondly, we consider global Poisson structures on the 4-dimensional sphere S^4 in Section 4. In general, to construct global Poisson structures, we solve the system of partial differential equations locally and attach the solutions globally. However it is too difficult, and we use another method. We regard S^4 as the 1-dimensional quaternionic projective space $\mathbb{HP}^1$, on which $\mathbb{CP}^3$ is the Hopf fibration. We study Poisson structures on $\mathbb{HP}^1$ associated with holomorphic Poisson structures on $\mathbb{CP}^3$. Then we obtain a lot of Poisson structures on $\mathbb{HP}^1$ and write down some examples, explicitly. Furthermore, we provide a real structure on the space of holomorphic Poisson structures on $\mathbb{CP}^3$. Then we can show that

Theorem 1.1. *The space of Poisson structures on $\mathbb{HP}^1$ induced by holomorphic Poisson structures on $\mathbb{CP}^3$ is the real form of the space of holomorphic Poisson structures on $\mathbb{CP}^3$ and it is a real algebraic variety.*

2 Preliminaries

In this section, we explain the definition of Poisson structure. For much of this material, we refer to [4]. Let M be a differentiable manifold of dimension n. A global section of $\wedge^q TM$ is called an *q-vector* on M. We write a 2-vector θ on M by $\theta = \sum_{i,j=1}^n \theta^{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}$ on a local coordinate $(x_1, \ldots, x_n)$ of M. The Schouten bracket $[\theta, \eta]_{Sch}$ for two 2-vectors θ, η is locally defined by

$$\begin{aligned}
[\theta,\eta]_{Sch} &= \sum_{i,j,k,l} [\theta^{ij}\frac{\partial}{\partial x_i}\wedge\frac{\partial}{\partial x_j}, \eta^{kl}\frac{\partial}{\partial x_k}\wedge\frac{\partial}{\partial x_l}]_{Sch} \\
&= \sum_{i,j,k,l} \Big\{ \theta^{ij}\frac{\partial\eta^{kl}}{\partial x_i}(\frac{\partial}{\partial x_j}\wedge\frac{\partial}{\partial x_k}\wedge\frac{\partial}{\partial x_l}) - \theta^{ij}\frac{\partial\eta^{kl}}{\partial x_j}(\frac{\partial}{\partial x_i}\wedge\frac{\partial}{\partial x_k}\wedge\frac{\partial}{\partial x_l}) \\
&\quad + \eta^{kl}\frac{\partial\theta^{ij}}{\partial x_k}(\frac{\partial}{\partial x_i}\wedge\frac{\partial}{\partial x_j}\wedge\frac{\partial}{\partial x_l}) - \eta^{kl}\frac{\partial\theta^{ij}}{\partial x_l}(\frac{\partial}{\partial x_i}\wedge\frac{\partial}{\partial x_j}\wedge\frac{\partial}{\partial x_k}) \Big\}
\end{aligned}$$

where $\theta = \sum_{i,j=1}^n \theta^{ij} \frac{\partial}{\partial x_i} \wedge \frac{\partial}{\partial x_j}$ and $\eta = \sum_{k,l=1}^n \eta^{kl} \frac{\partial}{\partial x_k} \wedge \frac{\partial}{\partial x_l}$.

We prepare a torsion free connection ∇ of M. Then the Schouten bracket is also globally defined by the following map

$$\begin{array}{ccccc}
\Gamma(\wedge^2 TM \otimes \wedge^2 TM) & \longrightarrow & \Gamma(\wedge^2 TM \otimes TM) & \longrightarrow & \Gamma(\wedge^3 TM) \\
\cup & & \cup & & \cup \\
(\theta, \eta) & \longmapsto & \nabla\theta\cdot\eta + \nabla\eta\cdot\theta & \longmapsto & \wedge^3(\nabla\theta\cdot\eta + \nabla\eta\cdot\theta)
\end{array}$$

where $\nabla\theta \cdot \eta$ is the convolution of $\nabla\theta \in \Gamma(\wedge^2 TM \otimes T^*M)$ and $\eta \in \Gamma(\wedge^2 TM)$ and the last map $\wedge^3$ means the antisymmetrization.

We call that a 2-vector θ is a *Poisson structure* on M if the Schouten bracket $[\theta, \theta]_{Sch}$ vanishes. The equation $[\theta, \theta]_{Sch} = 0$ is written as a system of partial differential equations that is so hard to solve. When the manifold is a complex manifold, we extend the Schouten bracket to complex 2-vectors by $\mathbb{C}$-linear. Then the

Schouten bracket $[\theta, \eta]_{Sch}$ for two holomorphic 2-vector θ, η is a holomorphic section of $\wedge^3 T^{1.0}M$. A holomorphic 2-vector θ is called a *holomorphic Poisson structure* if the Schouten bracket $[\theta, \theta]_{Sch}$ vanishes. A holomorphic Poisson structure θ induces two (real) Poisson structures as its real part and imaginary part. On a complex two dimensional manifold, any holomorphic 2-vector is a holomorphic Poisson structure. However, it is even hard to find a holomorphic Poisson structure in the case that the complex dimension is greater than and equal to three.

3 Holomorphic Poisson structures on $\mathbb{CP}^3$

3.1 Holomorphic 1-vectors on $\mathbb{CP}^3$

We denote by (z_0, z_1, z_2, z_3) the standard coordinate of $\mathbb{C}^4$. Let π be the projection $\mathbb{C}^4\backslash\{0\} \to \mathbb{CP}^3$. Any holomorphic 1-vector on $\mathbb{CP}^3$ is represented as the pushed foreword of a holomorphic 1-vector $\sum_{i,j} a_{ij} z_i \frac{\partial}{\partial z_j}$ on $\mathbb{C}^4\backslash\{0\}$ whose coefficients are polynomials of degree 1. We consider the correspondence given by

$$\sum_{i,j} a_{ij} z_i \frac{\partial}{\partial z_j} \mapsto (a_{ij}). \tag{1}$$

Then we obtain the isomorphism

$$\Big\{ \sum_{i,j} a_{ij} z_i \frac{\partial}{\partial z_j} \;\Big|\; a_{ij} \in \mathbb{C} \Big\} \cong gl(4, \mathbb{C})$$

where $gl(4, \mathbb{C})$ is the Lie algebra of the general linear group $\mathrm{GL}(4, \mathbb{C})$ of $\mathbb{C}^4$. The action of $\mathbb{C}^*$ on $\mathbb{C}^4\backslash\{0\}$ induces the vector field

$$l = z_0 \frac{\partial}{\partial z_0} + z_1 \frac{\partial}{\partial z_1} + z_2 \frac{\partial}{\partial z_2} + z_3 \frac{\partial}{\partial z_3}$$

which corresponds to the unit matrix E by the map (1). Then the 1-dimensional space $\langle\, l \,\rangle_{\mathbb{C}}$ is the space of holomorphic 1-vectors on $\mathbb{C}^4\backslash\{0\}$ which vanish by the push foreword $d\pi$. Hence, the space of holomorphic 1-vectors on $\mathbb{CP}^3$ can be identified with the quotient space

$$\Big\{ \sum_{i,j} a_{ij} z_i \frac{\partial}{\partial z_j} \;\Big|\; a_{ij} \in \mathbb{C} \Big\} \,/\, \langle\, l \,\rangle_{\mathbb{C}}$$

which is isomorphic to

$$gl(4, \mathbb{C}) / \langle E \rangle_{\mathbb{C}} \cong sl(4, \mathbb{C})$$

where $sl(4, \mathbb{C})$ is the Lie algebra of the special linear group $\mathrm{SL}(4, \mathbb{C})$.

3.2 Holomorphic 2-vectors on $\mathbb{CP}^3$

A holomorphic 2-vector on $\mathbb{CP}^3$ is the pushed foreword of a holomorphic 2-vector on $\mathbb{C}^4\backslash\{0\}$ whose coefficients are polynomials of degree 2. Such a 2-vector can be represented by $a_{ijkl}z_iz_j\frac{\partial}{\partial z_k}\wedge\frac{\partial}{\partial z_l}$ for a complex number a_{ijkl}. Then we can consider the coefficients a_{ijkl} as the element (a_{ijkl}) of the tensor space $S^2\mathbb{C}^4\otimes\wedge^2(\mathbb{C}^4)^\vee$ where $S^2\mathbb{C}^4$ is the symmetric 2-tensor space of $\mathbb{C}^4$ and $\wedge^2(\mathbb{C}^4)^\vee$ is the anti-symmetric 2-tensor space of the dual space $(\mathbb{C}^4)^\vee$ of $\mathbb{C}^4$. Any holomorphic 2 vector which vanishes by $d\pi$ is represented by $(\sum_{i,j}a_{ij}z_i\frac{\partial}{\partial z_j})\wedge l$ on $\mathbb{C}^4\backslash\{0\}$. Hence we can regard the space of holomorphic 2-vectors on $\mathbb{CP}^3$ as the quotient space

$$\Big\{\sum_{i,j,k,l}a_{ijkl}z_iz_j\frac{\partial}{\partial z_k}\wedge\frac{\partial}{\partial z_l}\;\Big|\;a_{ijkl}\in\mathbb{C}\Big\}\Big/\Big\{(\sum_{i,j}a_{ij}z_i\frac{\partial}{\partial z_j})\wedge l\;\Big|\;a_{ij}\in\mathbb{C}\Big\}$$

which is identified with

$$S^2\mathbb{C}^4\otimes\wedge^2(\mathbb{C}^4)^\vee\;/\;S^2\otimes\wedge^2(sl(4,\mathbb{C})\otimes\langle E\rangle_{\mathbb{C}})$$

where $S^2\otimes\wedge^2$ is the projection from $gl(\mathbb{C}^4)\otimes gl(\mathbb{C}^4)\cong(\otimes^2\mathbb{C}^4)\otimes(\otimes^2(\mathbb{C}^4)^\vee)$ to $S^2\mathbb{C}^4\otimes\wedge^2(\mathbb{C}^4)^\vee$.

As the same manner, the space of $\mathbb{C}^*$-invariant 3-vectors on $\mathbb{C}^4\backslash\{0\}$ is identified with

$$S^3\mathbb{C}^4\otimes\wedge^3(\mathbb{C}^4)^\vee$$

where $S^3\mathbb{C}^4$ is the symmetric 3-tensor space of $\mathbb{C}^4$ and $\wedge^3(\mathbb{C}^4)^\vee$ is the anti-symmetric 3-tensor space of $(\mathbb{C}^4)^\vee$. Then, the space of push forewords of $\mathbb{C}^*$-invariant 3-vectors on $\mathbb{C}^4\backslash\{0\}$ is isomorphic to

$$S^3\mathbb{C}^4\otimes\wedge^3(\mathbb{C}^4)^\vee\,/\,S^3\otimes\wedge^3(\otimes^2sl(4,\mathbb{C})\otimes\langle E\rangle_{\mathbb{C}})$$

where $S^3\otimes\wedge^3$ is the projection from $\otimes^3gl(\mathbb{C}^4)\cong(\otimes^3\mathbb{C}^4)\otimes(\otimes^3(\mathbb{C}^4)^\vee)$ to $S^3\mathbb{C}^4\otimes\wedge^3(\mathbb{C}^4)^\vee$.

3.3 The Schouten bracket on $\mathbb{CP}^3$

Let (a_{ijkl}) be an element of $S^2\mathbb{C}^4\otimes\wedge^2(\mathbb{C}^4)^\vee$. By fixing j and l, we can regard (a_{ijkl}) as the matrix whose (i,k)-component is a_{ijkl}, and denote the matrix by A_{jl}. Then, the matrix (A_{jl}) is the $gl(\mathbb{C}^4)$-valued matrix with the (j,l)-component A_{jl}. Thus, (a_{ijkl}) is associated with the $gl(\mathbb{C}^4)$-valued matrix (A_{jl}). If we take $(a_{ijkl}),(a'_{i'j'k'l'})\in S^2\mathbb{C}^4\otimes\wedge^2(\mathbb{C}^4)^\vee$ and denote by (A_{jl}) and $(A'_{j'l'})$ the associated matrices, respectively. Then we consider the bracket $[A_{jl},A'_{j'l'}]=A_{jl}A'_{j'l'}-A'_{j'l'}A_{jl}$ of matrices $A_{jl},A'_{j'l'}$ for each $j,l,j'l'$ and have the $gl(\mathbb{C}^4)$-valued matrix $([A_{jl},A'_{j'l'}])$ whose (j,l)-component is $[A_{jl},A'_{j'l'}]$. Moreover, $(([A_{jl},A'_{j'l'}]))$ is the $\otimes^2gl(\mathbb{C}^4)$-valued matrix whose (j',l')-component is $([A_{jl},A'_{j'l'}])$. We denote such $(([A_{jl},A'_{j'l'}]))$ by $[A_{jl},A'_{j'l'}]$ for simplicity. Thus, we can determine the element $[A_{jl},A'_{j'l'}]$ of $\otimes^3gl(\mathbb{C}^4)$ for $(a_{ijkl}),(a'_{i'j'k'l'})\in S^2\mathbb{C}^4\otimes\wedge^2(\mathbb{C}^4)^\vee$.

Under the identification of the space of $\mathbb{C}^*$-invariant q-vectors on $\mathbb{C}^4\backslash\{0\}$ with $S^q\mathbb{C}^4 \otimes \wedge^q(\mathbb{C}^4)^\vee$, the Schouten bracket for $\mathbb{C}^*$-invariant 2-vectors on $\mathbb{C}^4\backslash\{0\}$ can be considered as the map

$$\widetilde{F} : S^2\mathbb{C}^4 \otimes \wedge^2(\mathbb{C}^4)^\vee \times S^2\mathbb{C}^4 \otimes \wedge^2(\mathbb{C}^4)^\vee \to S^3\mathbb{C}^4 \otimes \wedge^3(\mathbb{C}^4)^\vee$$

defined by

$$\widetilde{F}(a_{ijkl}, a'_{i'j'k'l'}) = S^3 \otimes \wedge^3(4[A_{jl}, A'_{j'l'}])$$

for $a_{ijkl}, a'_{i'j'k'l'} \in S^2\mathbb{C}^4 \otimes \wedge^2(\mathbb{C}^4)^\vee$. Them, $\widetilde{F}$ is quadratic. Let u, v, u', v' be $\mathbb{C}^*$-invariant 1-vectors on $\mathbb{C}^4\backslash\{0\}$. Then, we obtain

$$[d\pi(w), d\pi(w')]_{Sch} = d\pi[w, w']_{Sch} \tag{2}$$

for $\mathbb{C}^*$-invariant 2-vectors w, w' on $\mathbb{C}^4\backslash\{0\}$. It follows from the π-relation (2) that the Schouten bracket on $\mathbb{CP}^3$ is given by the map from $S^2\mathbb{C}^4 \otimes \wedge^2(\mathbb{C}^4)^\vee/S^2 \otimes \wedge^2(sl(4,\mathbb{C})\otimes\langle E\rangle_{\mathbb{C}}) \times S^2\mathbb{C}^4\otimes\wedge^2(\mathbb{C}^4)^\vee/S^2\otimes\wedge^2(sl(4,\mathbb{C})\otimes\langle E\rangle_{\mathbb{C}})$ to $S^3\mathbb{C}^4\otimes\wedge^3(\mathbb{C}^4)^\vee/S^3\otimes \wedge^3(\otimes^2 sl(4,\mathbb{C}) \otimes \langle E\rangle_{\mathbb{C}})$ induced by $\widetilde{F}$. We denote by F the induced map. Then, we have

Proposition 3.1. *The space of holomorphic Poisson structures on $\mathbb{CP}^3$ is identified with*

$$\{a \in S^2\mathbb{C}^4 \otimes \wedge^2(\mathbb{C}^4)^\vee \,/\, S^2 \otimes \wedge^2(sl(4,\mathbb{C}) \otimes \langle E\rangle_{\mathbb{C}}) \mid F(a,a) = 0\}.$$

□

3.4 The Schouten bracket on local coordinates

We start to represent a holomorphic 2-vector $w = \sum_{i,j,k,l} a_{ijkl} z_i z_j \frac{\partial}{\partial z_k} \wedge \frac{\partial}{\partial z_l}$ on $\mathbb{C}^4\backslash\{0\}$ by using the local coordinate of $\mathbb{CP}^3$. Let U_r be the open set $\{[z_0, z_1, z_2, z_3] \in \mathbb{CP}^3 \mid z_r \neq 0\}$ of $\mathbb{CP}^3$ for $r = 0, 1, 2, 3$. We denote by ζ_i the function $\zeta_i = \frac{z_i}{z_r}$ on U_r for each $i = 0, 1, 2, 3$. Then the system of functions $\{\zeta_0, \zeta_1, \zeta_2, \zeta_3\}\backslash\{\zeta_r\}$ without ζ_r defines the inhomogeneous coordinate on U_r. We remark that $\zeta_r = 1$ on U_r. The 2-vector $d\pi(w)$ is represented by

$$d\pi(w) = \sum_{i,j,k\neq r, l\neq r} (a_{ijkl}\zeta_i\zeta_j - a_{ijkr}\zeta_i\zeta_j\zeta_k - a_{ijrl}\zeta_i\zeta_j\zeta_l)\frac{\partial}{\partial\zeta_k} \wedge \frac{\partial}{\partial\zeta_l} \tag{3}$$

on the local coordinate U_r. For simplicity, we denote by A_{rkl} the coefficient of $\frac{\partial}{\partial\zeta_k} \wedge \frac{\partial}{\partial\zeta_l}$ in (3) :

$$A_{rkl} = \sum_{i,j\neq r} a_{ijkl}\zeta_i\zeta_j - a_{ijkr}\zeta_i\zeta_j\zeta_k - a_{ijrl}\zeta_i\zeta_j\zeta_l.$$

We represent the Schouten bracket of $d\pi(w)$ on the local coordinates of $\mathbb{CP}^3$. For each r, we can take a triple (a, b, c) of $a, b, c, \in \{0, 1, 2, 3\}\backslash\{r\}$ with $a < b < c$. Then

the 3-vector $\frac{\partial}{\partial\zeta_a}\wedge\frac{\partial}{\partial\zeta_b}\wedge\frac{\partial}{\partial\zeta_c}$ generates the set of holomorphic 3-vectors on U_r, and it is denoted by w_r. Then the Schouten bracket $[d\pi(w), d\pi(w)]_{Sch}$ is written by

$$[d\pi(w), d\pi(w)]_{Sch} = 2\sum_{(k,l,m)\in R_r}(\frac{\partial A_{rkl}}{\partial\zeta_l}A_{rlm} - A_{rkl}\frac{\partial A_{rlm}}{\partial\zeta_l})w_r$$

on each U_r, where R_r be the set {(a,b,c), (b,c,a), (c,a,b)} of the rotation of the triple (a,b,c). Hence the Schouten bracket $[d\pi(w), d\pi(w)]_{Sch}$ vanishes if and only if

$$\sum_{(k,l,m)\in R_r}(\frac{\partial A_{rkl}}{\partial\zeta_l}A_{rlm} - A_{rkl}\frac{\partial A_{rlm}}{\partial\zeta_l}) = 0$$

on each U_r. It turns out that the space of holomorphic Poisson structures on $\mathbb{CP}^3$ is an algebraic variety. The space has the $\mathbb{C}^*$-action induced by the constant multiplication of 2-vectors. Then we obtain the following :

Proposition 3.2. *The quotient of the space of holomorphic Poisson structures on $\mathbb{CP}^3$ by the $\mathbb{C}^*$-action is a projective algebraic variety.* □

3.5 Examples of Poisson structures on $\mathbb{CP}^3$

In this section, we provide examples of Poisson structures on $\mathbb{CP}^3$. The Schouten bracket of the wedge product $X\wedge Y$ for two 1-vectors X and Y is given by

$$[X\wedge Y, X\wedge Y]_{Sch} = 2(\nabla_X Y - \nabla_Y X)\wedge X\wedge Y = 2[X,Y]\wedge X\wedge Y$$

where $[X,Y]$ means the Lie bracket for vector fields X and Y. It implies that $[X\wedge Y, X\wedge Y]_{Sch} = 0$ is the integrability condition of the distribution spanned by X and Y. Especially, for two 1-vectors X and Y with $[X,Y]=0$, the 2-vector $X\wedge Y$ is a Poisson structure. Let X and Y be 1-vectors on $\mathbb{CP}^3$ such that $X = d\pi(u), Y = d\pi(v)$ for 1-vectors u, v on $\mathbb{C}^4\backslash\{0\}$. Then $X\wedge Y$ is a Poisson structure if $[u,v]=0$ since $[X,Y] = d\pi[u,v]$. Thus we have a lot of examples of Poisson structures.

Example 3.3. We define w_1 by the 2-vector

$$w_1 = z_0 z_2\frac{\partial}{\partial z_1}\wedge\frac{\partial}{\partial z_3}$$

on $\mathbb{C}^4\backslash\{0\}$. Then w_1 induces a holomorphic Poisson structure $d\pi(w_1)$ on $\mathbb{CP}^3$. Actually, the Schouten bracket of $d\pi(w_1)$ is zero by the above argument and $d\pi(w_1)$ is a non-zero 2-vector on $\mathbb{CP}^3$ since it is represented as $\zeta_2\frac{\partial}{\partial\zeta_1}\wedge\frac{\partial}{\partial\zeta_3}$ on U_0. Hence $d\pi(w_1)$ is a non-trivial Poisson structure on $\mathbb{CP}^3$.

Example 3.4. We take the 2-vector

$$w_2 = (z_0\frac{\partial}{\partial z_1} - z_1\frac{\partial}{\partial z_0})\wedge(z_2\frac{\partial}{\partial z_3} - z_3\frac{\partial}{\partial z_2})$$

on $\mathbb{C}^4\backslash\{0\}$. Then the Schouten bracket of $d\pi(w_2)$ vanishes on $\mathbb{CP}^3$. The 2-vector $d\pi(w_2)$ is represented by $-\zeta_3(1+(\zeta_1)^2)\frac{\partial}{\partial\zeta_1}\wedge\frac{\partial}{\partial\zeta_2} - \zeta_2(1+(\zeta_1)^2)\frac{\partial}{\partial\zeta_3}\wedge\frac{\partial}{\partial\zeta_1} + \zeta_1((\zeta_2)^2 + (\zeta_3)^2)\frac{\partial}{\partial\zeta_2}\wedge\frac{\partial}{\partial\zeta_3}$ on U_0. Thus $d\pi(w_2)$ is a Poisson structure on $\mathbb{CP}^3$.

Example 3.5. We define w_3 by the 2-vector

$$w_3 = (z_0\frac{\partial}{\partial z_1} + z_1\frac{\partial}{\partial z_0} + z_2\frac{\partial}{\partial z_3} + z_3\frac{\partial}{\partial z_2}) \wedge (z_0\frac{\partial}{\partial z_3} + z_1\frac{\partial}{\partial z_2} + z_2\frac{\partial}{\partial z_1} + z_3\frac{\partial}{\partial z_0})$$

on $\mathbb{C}^4\backslash\{0\}$. Then the push foreword $d\pi(w_3)$ is a non-zero 2-vector on $\mathbb{CP}^3$ since the 0-degree part of the coefficient in (3) is a non-zero constant $a_{0013}\zeta_0\zeta_0 = 1$ on U_0. Hence $d\pi(w_3)$ is a Poisson structure on $\mathbb{CP}^3$.

If we take two 2-vectors $X \wedge Y$ and $X' \wedge Y'$ for 1-vectors X, Y, X', Y' on $\mathbb{CP}^3$, then we obtain

$$\begin{aligned}[X \wedge Y, X' \wedge Y']_{Sch} &= [X, X'] \wedge Y \wedge Y' - [X, Y'] \wedge Y \wedge X' \\ &\quad -[Y, X'] \wedge X \wedge Y' + [Y, Y'] \wedge X \wedge X'.\end{aligned}$$

Hence the 2-vector $X \wedge Y + X' \wedge Y'$ is a Poisson structure on $\mathbb{CP}^3$ if $[X, Y] = [X, X'] = [X, Y'] = [Y, X'] = [Y, Y'] = [X', Y'] = 0$. In general, if we take n 1-vectors $X_1, \ldots, X_n$ on $\mathbb{CP}^3$ and consider the 2-vector $w = \sum_{i<j} a_{ij} X_i \wedge X_j$ for a constant $a_{ij} \in \mathbb{C}$. Then w is a Poisson structure if $[X_i, X_j] = 0$ for $i, j = 1, \ldots, n$ since $[w, w]_{Sch}$ is generated by $[X_i, X_j] \wedge X_k \wedge X_l$ for $i, j, k, l = 1, \ldots, n$. In the case that each 1-vector X_i can be represented by $d\pi(u_i)$ of a 1-vector u_i on $\mathbb{C}^4\backslash\{0\}$, the 2-vector $w = \sum_{i<j} a_{ij} X_i \wedge X_j$ is a Poisson structure if $[u_i, u_j] = 0$ for $i, j = 1, \ldots, n$ (c.f. Theorem 5.3 and the equation (5.10) in [2]).

Example 3.6. We define u_1, u_2, u_3 by the three 1-vectors

$$u_1 = z_1\frac{\partial}{\partial z_1},\ u_2 = z_2\frac{\partial}{\partial z_2},\ u_3 = z_3\frac{\partial}{\partial z_3}$$

on $\mathbb{C}^4\backslash\{0\}$. Then, it is easy to see that $[u_i, u_j] = 0$ and $d\pi(u_i \wedge u_j) \neq 0$ for $i, j = 1, 2, 3$. Hence, we obtain a family of the Poisson structures

$$a_1 d\pi(u_1 \wedge u_2) + a_2 d\pi(u_1 \wedge u_3) + a_3 d\pi(u_2 \wedge u_3)$$

for $a_1, a_2, a_3 \in \mathbb{C}$.

It is known that there exist rich holomorphic foliations of codimension 1 on $\mathbb{CP}^3$. Especially, holomorphic foliations of degree 2 are completely classified [1]. Such a foliation is given by a holomorphic 1-form ω on $\mathbb{C}^4\backslash\{0\}$. The 1-form ω induces a 2-vector w by the correspondence

$$\omega = i_{l \wedge w}(dz_0 \wedge dz_1 \wedge dz_2 \wedge dz_3)$$

on $\mathbb{C}^4\backslash\{0\}$. Then, the push foreword $d\pi(w)$ is the Poisson structure on $\mathbb{CP}^3$. Hence, we can construct many Poisson structures corresponding to holomorphic foliations on $\mathbb{CP}^3$. We remark that w is not uniquely determined by the 1-form ω since there exists an ambiguity for 2-vectors factorized by l.

Example 3.7. We provide the 1-form ω by

$$\begin{aligned}\omega &= z_3(2z_1^2-3z_0z_2)dz_0+z_3(3z_2z_3-z_0z_1)dz_1\\ &\quad +z_3(z_0^2-2z_1z_3)dz_2-(z_0z_1^2-2z_0^2z_2+z_1z_2z_3)dz_3\end{aligned}$$

on $\mathbb{C}^4\backslash\{0\}$. Then we obtain the associated 2-vector

$$w=(z_0^2-2z_1z_3)\frac{\partial}{\partial z_0}\wedge\frac{\partial}{\partial z_1}+(z_0z_1-3z_2z_3)\frac{\partial}{\partial z_0}\wedge\frac{\partial}{\partial z_2}+(2z_1^2-3z_0z_2)\frac{\partial}{\partial z_1}\wedge\frac{\partial}{\partial z_2}$$

such that $d\pi(w)$ is the Poisson structure on $\mathbb{CP}^3$.

Example 3.8. When the 1-form ω' is given by

$$\omega'=\sum_{i,j=0}^{3}z_i(z_i^2-z_j^2)dz_j,$$

then the corresponding 2-vector

$$\begin{aligned}w' &= (z_3^2-z_2^2)\frac{\partial}{\partial z_0}\wedge\frac{\partial}{\partial z_1}+(z_1^2-z_3^2)\frac{\partial}{\partial z_0}\wedge\frac{\partial}{\partial z_2}+(z_2^2-z_1^2)\frac{\partial}{\partial z_0}\wedge\frac{\partial}{\partial z_3}\\ &\quad +(z_3^2-z_0^2)\frac{\partial}{\partial z_1}\wedge\frac{\partial}{\partial z_2}+(z_0^2-z_2^2)\frac{\partial}{\partial z_1}\wedge\frac{\partial}{\partial z_3}+(z_1^2-z_0^2)\frac{\partial}{\partial z_2}\wedge\frac{\partial}{\partial z_3}\end{aligned}$$

induces the Poisson structure $d\pi(w')$ on $\mathbb{CP}^3$.

4 Poisson structures on $\mathbb{HP}^1$

Let $\mathbb{H}$ be the Hamilton's quaternionic number field $\mathbb{R}+i\mathbb{R}+j\mathbb{R}+k\mathbb{R}$. Then $\mathbb{H}^2$ is isomorphic to the right $\mathbb{C}$-module $\mathbb{C}^4$ under the identification of $\mathbb{H}$ with $\mathbb{C}^2(\cong\mathbb{C}+j\mathbb{C})$. Let p be the projection from $\mathbb{C}^4\backslash\{0\}\cong\mathbb{H}^2\backslash\{0\}$ to $\mathbb{HP}^1$. In this section, we consider real q-vectors on $\mathbb{HP}^1$ given by the push foreword of $\mathbb{H}^*$-invariant real q-vectors with holomorphic $(q,0)$-parts on $\mathbb{C}^4\backslash\{0\}$.

4.1 $\mathbb{H}^*$-invariant real q-vectors on $\mathbb{C}^4\backslash\{0\}$

Let $\widetilde{V}_q$ be the space of $\mathbb{H}^*$-invariant real q-vectors with holomorphic $(q,0)$-parts on $\mathbb{C}^4\backslash\{0\}$:

$$\widetilde{V}_q=\{v\in\Gamma_{\mathbb{H}^*\text{-inv.}}(\wedge^qT\mathbb{C}^4\backslash\{0\})\mid v^{q,0}:\text{holomorphic}\,\}.$$

The right action of j on $\mathbb{C}^4\backslash\{0\}$ induces the map $j:\mathbb{C}^4\backslash\{0\}\to\mathbb{C}^4\backslash\{0\}$ given by

$$(z_0,z_1,z_2,z_3)\mapsto(-\bar{z}_1,\bar{z}_0,-\bar{z}_3,\bar{z}_2)$$

for $(z_0,z_1,z_2,z_3)\in\mathbb{C}^4\backslash\{0\}$. Let l' denote the complex vector field on $\mathbb{C}^4\backslash\{0\}$ generated by $\exp(tj)$ for $t\in\mathbb{R}$. Then we have

$$l'=\bar{z}_0\frac{\partial}{\partial z_1}-\bar{z}_1\frac{\partial}{\partial z_0}+\bar{z}_2\frac{\partial}{\partial z_3}-\bar{z}_3\frac{\partial}{\partial z_2}.$$

Any $\mathbb{C}^*$-invariant real q-vector on $\mathbb{C}^4\backslash\{0\}$ is $\mathbb{H}^*$-invariant if and only if it is preserved by l'. Hence, the space $\widetilde{V}_q$ is written by

$$\widetilde{V}_q = \{v \in \Gamma_{\mathbb{C}^*\text{-inv.}}(\wedge^q T\mathbb{C}^4\backslash\{0\}) \mid L_{l'}v = 0,\ v^{q,0} : \text{holomorphic}\}$$

where $L_{l'}v$ means the Lie derivative of v by l'. We also define V_q by the space of real q-vectors on $\mathbb{HP}^1$ induced by elements of $\widetilde{V}_q$:

$$V_q = \{dp(v) \in \Gamma(\wedge^q T\mathbb{HP}^1) \mid v \in \widetilde{V}_q\}.$$

The real and imaginary parts of l and l' span the real 4-dimensional vector space. Then the intersection of the space and $\widetilde{V}_1$ is the 1-dimensional vector space generated by $l + \bar{l}$. Thus the space $\langle l + \bar{l}\rangle_{\mathbb{R}}$ is the kernel of the push foreword by p, and V_1 is identified with the the quotient space $\widetilde{V}_1/\langle l + \bar{l}\rangle_{\mathbb{R}}$. Moreover, there exists an isomorphism

$$V_q \cong \widetilde{V}_q \,/\, \widetilde{V}_{q-1} \wedge \langle l + \bar{l}\rangle_{\mathbb{R}}$$

where the right hand side is the quotient space of $\widetilde{V}_q$ divided by $\widetilde{V}_{q-1} \wedge \langle l + \bar{l}\rangle_{\mathbb{R}}$.

4.2 1-vectors on $\mathbb{HP}^1$

We consider the space $\widetilde{V}_1$ of $\mathbb{H}^*$-invariant real 1-vectors on $\mathbb{C}^4\backslash\{0\}$, whose $(1,0)$-parts are holomorphic. The space of $\mathbb{C}^*$-invariant real 1-vectors on $\mathbb{C}^4\backslash\{0\}$ with holomorphic $(1,0)$-parts is

$$\Big\{\sum_{i,j}(a_{ij}z_i\frac{\partial}{\partial z_j} + \bar{a}_{ij}\bar{z}_i\frac{\partial}{\partial \bar{z}_j}) \;\Big|\; a_{ij} \in \mathbb{C}\Big\}.$$

Let φ be the mapping $(0,1,2,3) \to (1,0,3,2)$, that is, $\varphi(0) = 1, \varphi(1) = 0, \varphi(2) = 3, \varphi(3) = 2$. Then, the 1-vector $\sum_{i,j}(a_{ij}z_i\frac{\partial}{\partial z_j} + \bar{a}_{ij}\bar{z}_i\frac{\partial}{\partial \bar{z}_j})$ is $\mathbb{H}^*$-invariant if and only if

$$a_{ij} = (-1)^{i+j}\bar{a}_{\varphi(i)\varphi(j)}. \tag{4}$$

Thus the space $\widetilde{V}_1$ is written by

$$\widetilde{V}_1 = \Big\{\sum_{i,j}(a_{ij}z_i\frac{\partial}{\partial z_j} + \bar{a}_{ij}\bar{z}_i\frac{\partial}{\partial \bar{z}_j}) \;\Big|\; a_{ij} = (-1)^{i+j}\bar{a}_{\varphi(i)\varphi(j)}\Big\}.$$

We can consider the coefficients a_{ij} as the matrix (a_{ij}) of $gl(4,\mathbb{C})$ under the correspondence

$$\sum_{i,j}(a_{ij}z_i\frac{\partial}{\partial z_j} + \bar{a}_{ij}\bar{z}_i\frac{\partial}{\partial \bar{z}_j}) \mapsto (a_{ij}).$$

Let $\mathrm{GL}(2,\mathbb{H})$ be the general linear group of $\mathbb{H}^2$. Then the Lie algebra $gl(2,\mathbb{H})$ of $\mathrm{GL}(2,\mathbb{H})$ is identified with the subspace of $gl(4,\mathbb{C})$ by the map

$$\begin{pmatrix} a_{00}+jb_{00} & a_{01}+jb_{01} \\ a_{10}+jb_{10} & a_{11}+jb_{11} \end{pmatrix} \longmapsto \begin{pmatrix} a_{00} & -\bar{b}_{00} & a_{01} & -\bar{b}_{01} \\ b_{00} & \bar{a}_{00} & b_{01} & \bar{a}_{01} \\ a_{10} & -\bar{b}_{10} & a_{11} & -\bar{b}_{11} \\ b_{10} & \bar{a}_{10} & b_{11} & \bar{a}_{11} \end{pmatrix} \tag{5}$$

for $(a_{kl} + jb_{kl}) \in gl(2, \mathbb{H})$. Conversely, an element (a_{ij}) of $gl(4, \mathbb{C})$ is in the image of $gl(2, \mathbb{H})$ by the map (5) if a_{ij} satisfies the equation (4). Hence, we obtain

$$\widetilde{V}_1 \cong gl(2, \mathbb{H}).$$

Since the real 1-vector $l + \bar{l}$ corresponds to the unit matrix $E \in gl(2, \mathbb{H})$, we also have

$$V_1 \cong gl(2, \mathbb{H}) / \langle E \rangle_{\mathbb{R}}.$$

4.3 2-vectors on $\mathbb{HP}^1$

We consider the space $\widetilde{V}_2$ of $\mathbb{H}^*$-invariant real 2-vectors on $\mathbb{C}^4 \backslash \{0\}$ with holomorphic $(2, 0)$-parts. We take a $\mathbb{C}^*$-invariant real 2-vector w on $\mathbb{C}^4 \backslash \{0\}$ given by

$$w = \sum_{i,j,k,l} a_{ijkl} z_i z_j \frac{\partial}{\partial z_k} \wedge \frac{\partial}{\partial z_l} + b_{ijkl} z_i \bar{z}_j \frac{\partial}{\partial z_k} \wedge \frac{\partial}{\partial \bar{z}_l} + \bar{a}_{ijkl} \bar{z}_i \bar{z}_j \frac{\partial}{\partial \bar{z}_k} \wedge \frac{\partial}{\partial \bar{z}_l}$$

for $a_{ijkl}, b_{ijkl} \in \mathbb{C}$. Then the equations $a_{jikl} = a_{ijkl},\ a_{ijlk} = -a_{ijkl},\ b_{jilk} = -\bar{b}_{ijkl}$ hold for $i, j, k, l = 0, \ldots, 3$. We recall that the coefficients a_{ijkl} define the element (a_{ijkl}) of $S^2\mathbb{C}^4 \otimes \wedge^2(\mathbb{C}^4)^\vee$ as in §3.2. The $\mathbb{H}^*$-invariant condition of w is equal to

$$b_{ijkl} = 2(-1)^{j+l} a_{i\varphi(j)k\varphi(l)} = 2(-1)^{i+k} \bar{a}_{\varphi(i)j\varphi(k)l}.$$

It yields that b_{ijkl} is uniquely determined by (a_{ijkl}) satisfying

$$a_{ijkl} = (-1)^{i+j+k+l} \bar{a}_{\varphi(i)\varphi(j)\varphi(k)\varphi(l)}. \tag{6}$$

The space of elements of $S^2\mathbb{C}^4 \otimes \wedge^2(\mathbb{C}^4)^\vee$ satisfying the condition (6) is the tensor space $S^2 \otimes \wedge^2(gl(2, \mathbb{H}) \otimes gl(2, \mathbb{H}))$. Hence, by the correspondence

$$\sum_{i,j,k,l} a_{ijkl} z_i z_j \frac{\partial}{\partial z_k} \wedge \frac{\partial}{\partial z_l} + b_{ijkl} z_i \bar{z}_j \frac{\partial}{\partial z_k} \wedge \frac{\partial}{\partial \bar{z}_l} + \bar{a}_{ijkl} \bar{z}_i \bar{z}_j \frac{\partial}{\partial \bar{z}_k} \wedge \frac{\partial}{\partial \bar{z}_l} \mapsto (a_{ijkl}),$$

we obtain the isomorphism

$$\widetilde{V}_2 \cong S^2 \otimes \wedge^2(gl(2, \mathbb{H}) \otimes gl(2, \mathbb{H})).$$

It yields that

$$V_2 \cong S^2 \otimes \wedge^2(gl(2, \mathbb{H}) \otimes gl(2, \mathbb{H})) \,/\, S^2 \otimes \wedge^2(gl(2, \mathbb{H}) \otimes \langle E \rangle_{\mathbb{R}}).$$

In the case of $q = 3$, we can find that the space $\widetilde{V}_3$ is identified with the tensor space $S^3 \otimes \wedge^3(\otimes^3 gl(2, \mathbb{H}))$:

$$\widetilde{V}_3 \cong S^3 \otimes \wedge^3(\otimes^3 gl(2, \mathbb{H})).$$

Then we also have the isomorphism

$$V_3 \cong S^3 \otimes \wedge^3(\otimes^3 gl(2, \mathbb{H})) \,/\, S^3 \otimes \wedge^3(\otimes^2 gl(2, \mathbb{H}) \otimes \langle E \rangle_{\mathbb{R}}).$$

We will see that the space V_q is identified with a real form of the space induced by $\mathbb{C}^*$-invariant holomorphic q-vectors on $\mathbb{C}^4 \backslash \{0\}$ in §4.5.

4.4 $\mathbb{C}^*$-invariant holomorphic q-vectors on $\mathbb{C}^4\backslash\{0\}$

Let $\widetilde{W}_q$ be the space of $\mathbb{C}^*$-invariant holomorphic q-vectors on $\mathbb{C}^4\backslash\{0\}$:

$$\widetilde{W}_q = \{w \in \Gamma_{\mathbb{C}^*\text{-inv.}}(\wedge^q T^{\mathbb{C}}\mathbb{C}^4\backslash\{0\}) \mid w : \text{holomorphic}\,\}.$$

We define W_q as the space of the push foreword of elements of $\widetilde{W}_q$ by the projection $\pi : \mathbb{C}^4\backslash\{0\} \to \mathbb{CP}^3$:

$$W_q = \{d\pi(w) \in \Gamma(\wedge^q T^{\mathbb{C}}\mathbb{CP}^3) \mid w \in \widetilde{W}_q\}.$$

Then the space W_q is isomorphic to the quotient space $\widetilde{W}_q \,/\, \widetilde{W}_{q-1}\wedge\langle l\rangle_{\mathbb{C}}$ of $\widetilde{W}_q$ divided by $\widetilde{W}_{q-1} \wedge \langle l\rangle_{\mathbb{C}}$. We recall that there exist isomorphisms

$$\begin{aligned}\widetilde{W}_q &\cong S^q\mathbb{C}^4 \otimes \wedge^q(\mathbb{C}^4)^\vee,\\ W_q &\cong S^q\mathbb{C}^4 \otimes \wedge^q(\mathbb{C}^4)^\vee \,/\, S^q \otimes \wedge^q(\otimes^{q-1} sl(4,\mathbb{C}) \otimes \langle E\rangle_{\mathbb{C}}).\end{aligned}$$

4.5 The real structures on $\widetilde{W}_q$ and W_q

We define the map $\Phi : \Gamma(T^{\mathbb{C}}\mathbb{C}^4\backslash\{0\}) \to \Gamma(T^{\mathbb{C}}\mathbb{C}^4\backslash\{0\})$ by

$$\Phi(u) = \overline{dj(u)}$$

for any $u \in \Gamma(T^{\mathbb{C}}\mathbb{C}^4\backslash\{0\})$. Then the map Φ is anti-$\mathbb{C}$-linear and preserves the inner product given by the standard Riemannian metric on $\mathbb{C}^4$. Since $\Phi^2(u) = u$ for any $\mathbb{C}^*$-invariant holomorphic 1-vector u on $\mathbb{C}^4\backslash\{0\}$, the map Φ defines the real structure on the space $\widetilde{W}_1$ of $\mathbb{C}^*$-invariant holomorphic 1-vectors on $\mathbb{C}^4\backslash\{0\}$. We denote by $\widetilde{W}_1^{\mathrm{Re}}$ the real form of Φ, that is, the set of fixed points of Φ. Then the space $\widetilde{V}_1$ of $\mathbb{H}^*$-invariant real 1-vectors on $\mathbb{C}^4\backslash\{0\}$ with holomorphic $(1,0)$-parts is isomorphic to $\widetilde{W}_1^{\mathrm{Re}}$ by the map $v \mapsto v^{1,0}$ for $v \in \widetilde{V}_1$. The map Φ induces the real structure on the space $\widetilde{W}_q$ of $\mathbb{C}^*$-invariant holomorphic q-vectors on $\mathbb{C}^4\backslash\{0\}$, which is denoted by Φ for simplicity. Then the space $\widetilde{V}_q$ is isomorphic to the real form $\widetilde{W}_q^{\mathrm{Re}}$ of Φ by the map

$$v \mapsto v^{q,0} \tag{7}$$

for $v \in \widetilde{V}_q$. Under the identification of $\widetilde{W}_q$ with $S^q\mathbb{C}^4 \otimes \wedge^q(\mathbb{C}^4)^\vee$, the real structure Φ can be written by

$$\Phi(a_{i_1 i_2 \cdots i_{2q-1} i_{2q}}) = (-1)^{i_1+i_2+\cdots+i_{2q}} \bar{a}_{\varphi(i_1)\varphi(i_2)\cdots\varphi(i_{2q})}$$

for $(a_{i_1 i_2 \cdots i_{2q-1} i_{2q}}) \in S^q\mathbb{C}^4 \otimes \wedge^q(\mathbb{C}^4)^\vee$. It follows from $\Phi(l) = l$ that Φ preserves the complex vector space $\langle l\rangle_{\mathbb{C}}$. Hence, Φ provides the real structure on the space $W_q \cong \widetilde{W}_q \,/\, \widetilde{W}_{q-1} \wedge \langle l\rangle_{\mathbb{C}}$. We denote by Φ the real structure on W_q, again. Then, we regard the space V_q as the real form W_q^{Re} of W_q since

$$V_q \cong \widetilde{V}_q \,/\, \widetilde{V}_{q-1} \wedge \left\langle l + \bar{l}\right\rangle_{\mathbb{R}} \cong \widetilde{W}_q^{\mathrm{Re}} \,/\, \widetilde{W}_{q-1}^{\mathrm{Re}} \wedge \langle l\rangle_{\mathbb{R}} \cong W_q^{\mathrm{Re}}$$

where the second isomorphism is given by the map (7).

4.6 The Schouten bracket on $\mathbb{HP}^1$

The Schouten bracket of $\mathbb{C}^4\backslash\{0\}$ satisfies the relation

$$[\Phi(w), \Phi(w')]_{Sch} = \Phi[w, w']_{Sch}$$

for any 2-vectors w, w' on $\mathbb{C}^4\backslash\{0\}$ since Φ is isometric and preserves the Riemannian connection on $\mathbb{C}^4\backslash\{0\}$. We recall that the map Φ defines the real structure on the space $\widetilde{W}_2$ of $\mathbb{C}^*$-invariant holomorphic 2-vectors on $\mathbb{C}^4\backslash\{0\}$. Let w, w' be two elements of the real form $\widetilde{W}_2^{\mathrm{Re}}$ of $\widetilde{W}_2$. Then the Schouten bracket $[w, w']_{Sch}$ is in the real form $\widetilde{W}_3^{\mathrm{Re}}$ of $\widetilde{W}_3$ since $[w, w']_{Sch} = [\Phi(w), \Phi(w')]_{Sch} = \Phi[w, w']_{Sch}$. Thus, the Schouten bracket maps $\widetilde{W}_2^{\mathrm{Re}} \times \widetilde{W}_2^{\mathrm{Re}}$ to $\widetilde{W}_3^{\mathrm{Re}}$.

If we take two elements u, v of $\widetilde{V}_2$, then the Schouten bracket $[u, v]_{Sch}$ is in $\widetilde{V}_3$. Actually, since $(2,0)$-parts $u^{2,0}, v^{2,0}$ are elements of $\widetilde{W}_2^{\mathrm{Re}}$, the $(3,0)$-part $[u, v]_{Sch}^{3,0} = [u^{2,0}, v^{2,0}]_{Sch}$ of $[u, v]_{Sch}$ is in $\widetilde{W}_3^{\mathrm{Re}}$. It means that $[u, v]_{Sch}$ is the element of $\widetilde{V}_3$. Hence, the Schouten bracket induces the map from $\widetilde{V}_2 \times \widetilde{V}_2$ to $\widetilde{V}_3$. Moreover, since $dp[u, v]_{Sch} = [dp(u), dp(v)]_{Sch}$ for $u, v \in \widetilde{V}_2$, it induces the map from $V_2 \times V_2$ to V_3, which is the Schouten bracket of $\mathbb{HP}^1$ restricted to V_2.

Under the identification of $\widetilde{W}_q$ with $S^q\mathbb{C}^4 \otimes \wedge^q(\mathbb{C}^4)^\vee$, the Schouten brackets on $\widetilde{W}_2$ and W_2 are given by the maps $\widetilde{F}$ and F as in §3.3, respectively. Then the Schouten bracket on V_2 corresponds the map F restricted to $S^2 \otimes \wedge^2(gl(2,\mathbb{H}) \otimes gl(2,\mathbb{H}))\,/\,S^2 \otimes \wedge^2(gl(2,\mathbb{H}) \otimes \langle E \rangle_{\mathbb{R}})$. Hence, we have

Theorem 4.1. *The space of Poisson structures on $\mathbb{HP}^1$ induced by holomorphic Poisson structures on $\mathbb{CP}^3$ is identified with*

$$\{a \in S^2 \otimes \wedge^2(gl(2,\mathbb{H}) \otimes gl(2,\mathbb{H}))\,/\,S^2 \otimes \wedge^2(gl(2,\mathbb{H}) \otimes \langle E \rangle_{\mathbb{R}}) \mid F(a,a) = 0\},$$

and it is the real form of the space of holomorphic Poisson structures on $\mathbb{CP}^3$. □

The space has the $\mathbb{R}^*$-action induced by the $\mathbb{C}^*$-action on the space of holomorphic Poisson structures on $\mathbb{CP}^3$. The quadratic polynomial $F(a,a)$ has real coefficients. Hence, we obtain

Corollary 4.2. *The quotient of the space of Poisson structures on $\mathbb{HP}^1$ induced by holomorphic Poisson structures on $\mathbb{CP}^3$ by the $\mathbb{R}^*$-action is a real projective algebraic variety.* □

4.7 Examples of Poisson structures on $\mathbb{HP}^1$

In order to check that $dp(w)$ is a non-zero 2-vector on $\mathbb{HP}^1$, we represent 1-vectors by using the local coordinate of $\mathbb{HP}^1$. Let V_m be the open set $\{[h_0, h_1] \in \mathbb{HP}^1 | h_m \neq 0\}$ of $\mathbb{HP}^1$ for $m = 0, 1$. The functions $s_0 = h_1 h_0^{-1}$ and $s_1 = h_0 h_1^{-1}$ provide coordinate on V_0 and V_1, respectively. We can write $h_0 = (z_0, z_1)$ and $h_1 = (z_2, z_3)$ by using the complex coordinate (z_0, z_1, z_2, z_3) of $\mathbb{C}^4\backslash\{0\}$. On the local coordinate (V_0, s_0), if we

denote the inhomogeneous coordinate s_0 by $t_0 + jt_1$, then we have

$$\begin{aligned} dp(\frac{\partial}{\partial z_0}) &= -t_0\frac{\partial}{\partial t_0} - t_1\frac{\partial}{\partial t_1}, \\ dp(\frac{\partial}{\partial z_1}) &= \bar{t}_1\frac{\partial}{\partial t_0} - \bar{t}_0\frac{\partial}{\partial t_1}, \\ dp(\frac{\partial}{\partial z_2}) &= \frac{\partial}{\partial t_0}, \\ dp(\frac{\partial}{\partial z_3}) &= \frac{\partial}{\partial t_1} \end{aligned}$$

at $z_0 = 1, z_1 = 0, z_2 = t_0, z_3 = t_1$. Similarly, on V_1 with the coordinate $s_1 = t_0 + jt_1$, we also have

$$\begin{aligned} dp(\frac{\partial}{\partial z_0}) &= \frac{\partial}{\partial t_0}, \\ dp(\frac{\partial}{\partial z_1}) &= \frac{\partial}{\partial t_1}, \\ dp(\frac{\partial}{\partial z_2}) &= -t_0\frac{\partial}{\partial t_0} - t_1\frac{\partial}{\partial t_1}, \\ dp(\frac{\partial}{\partial z_3}) &= \bar{t}_1\frac{\partial}{\partial t_0} - \bar{t}_0\frac{\partial}{\partial t_1} \end{aligned}$$

at $z_0 = t_0, z_1 = t_1, z_2 = 1, z_3 = 0$.

Let u and v be two complex 1-vectors on $\mathbb{C}^4\backslash\{0\}$. If $u + \bar{u}$ and $v + \bar{v}$ are $\mathbb{H}^*$-invariant, then the Schouten bracket of the real 2-vector $dp(u+\bar{u})\wedge dp(v+\bar{v})$ on $\mathbb{HP}^1$ is given by $2dp([u, v]+\overline{[u, v]})\wedge(u+\bar{u})\wedge(v+\bar{v}))$. It implies that $dp(u+\bar{u})\wedge dp(v+\bar{v})$ defines a Poisson structure in the case of $[u, v] = 0$. In general, if we take n 1-vectors $v_1, \ldots, v_n$ on $\mathbb{C}^4\backslash\{0\}$ such that $[v_i, v_j] = 0$ for $i, j = 1, \ldots, n$, then a real 2-vector $\sum_{i<j} a_{ij}dp(v_i + \bar{v}_i)\wedge dp(v_j + \bar{v}_j)$ is a Poisson structure for a constant $a_{ij} \in \mathbb{R}$ by the similar argument of $\mathbb{CP}^3$. We provide examples of Poisson structures of such type.

Example 4.3. We define 1-vectors v_1, v_2, v_3 by

$$v_1 = z_0\frac{\partial}{\partial z_0} + z_1\frac{\partial}{\partial z_1},\ v_2 = i(z_0\frac{\partial}{\partial z_0} - z_1\frac{\partial}{\partial z_1}),\ v_3 = i(z_2\frac{\partial}{\partial z_2} - z_3\frac{\partial}{\partial z_3})$$

on $\mathbb{C}^4\backslash\{0\}$. Then $[v_1, v_2] = [v_1, v_3] = [v_2, v_3] = 0$. We can also check that each push foreword $dp((v_1 + \bar{v}_1)\wedge(v_2 + \bar{v}_2)), dp((v_1 + \bar{v}_1)\wedge(v_3 + \bar{v}_3)), dp((v_2 + \bar{v}_2)\wedge(v_3 + \bar{v}_3))$ is a non-zero 2-vector on $\mathbb{HP}^1$ by using the local coordinate V_0. Hence we obtain a family of Poisson structures

$$a_1dp((v_1 + \bar{v}_1)\wedge(v_2 + \bar{v}_2)) + a_2dp((v_1 + \bar{v}_1)\wedge(v_3 + \bar{v}_3)) + a_3dp((v_2 + \bar{v}_2)\wedge(v_3 + \bar{v}_3))$$

for $a_1, a_2, a_3 \in \mathbb{R}$.

In the section 3.5, we consider a holomorphic 1-form ω on $\mathbb{C}^4\backslash\{0\}$ associated with a holomorphic foliation of degree 2 on $\mathbb{CP}^3$. The form ω induces the 2-vector w

such that $d\pi(w)$ is the Poisson structure on $\mathbb{CP}^3$. The coefficients of such a 1-form ω are polynomials of degree 3. We can define the real structure Φ on the space of 1-forms whose coefficients are polynomials of degree 3 as in the section 4.5. If the 2-vector w is real with respect to Φ, then the 1-form ω is so. However, the converse is not true by the choice of the 2-vector w for the 1-form ω. We provide an explicit correspondence from the 1-form ω to the 2-vector w as follows. We take a holomorphic 1-form ω on $\mathbb{C}^4\backslash\{0\}$ associated with a holomorphic foliation of degree 2 on $\mathbb{CP}^3$, and write the 1-form by

$$\omega = \sum_{i=0}^{3} f_i dz_i$$

where f_i is a polynomial of degree 3. Each polynomial f_i decomposes into

$$f_i = \sum_{j\neq i} f_{i,j}z_j + \sum_{\substack{j,k\neq i\\ j<k}} f_{i,jk}z_jz_k + \sum_{\substack{j,k,l\neq i\\ j<k<l}} f_{i,jkl}z_jz_kz_l$$

where $f_{i,j}$ is a polynomial of degree 2 generated by z_i and z_j, and $f_{i,jk}$ is a polynomial of degree 1 generated by z_i, z_j and z_k, and $f_{i,jkl}$ is a complex number. Then we obtain the 2-vector w corresponding to ω by

$$\begin{aligned}
w \;=\;& f_{0,1}\frac{\partial}{\partial z_2}\wedge\frac{\partial}{\partial z_3} + f_{2,0}\frac{\partial}{\partial z_1}\wedge\frac{\partial}{\partial z_3} + f_{0,3}\frac{\partial}{\partial z_1}\wedge\frac{\partial}{\partial z_2}\\
&+f_{1,2}\frac{\partial}{\partial z_0}\wedge\frac{\partial}{\partial z_3} + f_{3,1}\frac{\partial}{\partial z_0}\wedge\frac{\partial}{\partial z_2} + f_{2,3}\frac{\partial}{\partial z_0}\wedge\frac{\partial}{\partial z_1}\\
&+\frac{1}{3}\{((f_{3,12}-f_{2,13})z_1\frac{\partial}{\partial z_1} + (f_{1,23}-f_{3,12})z_2\frac{\partial}{\partial z_2} + (f_{2,13}-f_{1,23})z_3\frac{\partial}{\partial z_3})\wedge\frac{\partial}{\partial z_0}\\
&+((f_{2,03}-f_{3,02})z_0\frac{\partial}{\partial z_0} + (f_{3,02}-f_{0,23})z_2\frac{\partial}{\partial z_2} + (f_{0,23}-f_{2,03})z_3\frac{\partial}{\partial z_3})\wedge\frac{\partial}{\partial z_1}\\
&+((f_{3,01}-f_{1,03})z_0\frac{\partial}{\partial z_0} + (f_{0,13}-f_{3,01})z_1\frac{\partial}{\partial z_1} + (f_{1,03}-f_{0,13})z_3\frac{\partial}{\partial z_3})\wedge\frac{\partial}{\partial z_2}\\
&+((f_{1,02}-f_{2,01})z_0\frac{\partial}{\partial z_0} + (f_{2,01}-f_{0,12})z_1\frac{\partial}{\partial z_1} + (f_{0,12}-f_{1,02})z_2\frac{\partial}{\partial z_2})\wedge\frac{\partial}{\partial z_3}\}\\
&+\frac{1}{4}\{(f_{2,013}-f_{3,012})z_0z_1\frac{\partial}{\partial z_0}\wedge\frac{\partial}{\partial z_1} + (f_{3,012}-f_{1,023})z_0z_2\frac{\partial}{\partial z_0}\wedge\frac{\partial}{\partial z_2}\\
&+(f_{1,023}-f_{2,013})z_0z_3\frac{\partial}{\partial z_0}\wedge\frac{\partial}{\partial z_3} + (f_{0,123}-f_{3,012})z_1z_2\frac{\partial}{\partial z_1}\wedge\frac{\partial}{\partial z_2}\\
&+(f_{2,013}-f_{0,123})z_1z_3\frac{\partial}{\partial z_1}\wedge\frac{\partial}{\partial z_3} + (f_{0,123}-f_{1,023})z_2z_3\frac{\partial}{\partial z_2}\wedge\frac{\partial}{\partial z_3}\}
\end{aligned} \tag{8}$$

on $\mathbb{C}^4\backslash\{0\}$. Under the correspondence (8), the 2-vector w is real if ω is so. In [1], the space of holomorphic foliations of degree 2 on $\mathbb{CP}^3$ is decomposed into the six component. the component $\overline{R(2,2)}$ is consisted by 1-forms

$$gdf - fdg$$

for polynomials f, g of degree 2 on $\mathbb{C}^4\backslash\{0\}$. The space of polynomials of degree 2 on $\mathbb{C}^4\backslash\{0\}$ is isomorphic to $S^2\mathbb{C}^4$. If we take f, g in $gl(2,\mathbb{H})\cap S^2\mathbb{C}^4$, then $\omega = gdf - fdg$ is real with respect to Φ, and the corresponding 2-vector w defines the Poisson structure on $\mathbb{HP}^1$.

Example 4.4. We take the polynomials

$$f = z_0^2 + z_1^2, \quad g = \sqrt{-1}\, z_0 z_1$$

on $\mathbb{C}^4\backslash\{0\}$. The corresponding 2-vector

$$w = \sqrt{-1}\,(z_3^2 - z_2^2)\frac{\partial}{\partial z_0} \wedge \frac{\partial}{\partial z_1}$$

induces the Poisson structure $d\pi(w)$ on $\mathbb{CP}^3$. The 2-vector w is real since $f, g \in gl(2,\mathbb{H})\cap S^2\mathbb{C}^4$. Hence, the 2-vector w defines the Poisson structure on $\mathbb{HP}^1$.

Example 4.5. For the polynomials

$$f = z_0^2 + z_1^2 + z_2^2 + z_3^2, \quad g = \sqrt{-1}\, z_0 z_1$$

on $\mathbb{C}^4\backslash\{0\}$, the corresponding 2-vector is given by

$$\begin{aligned} w \;=\; & \sqrt{-1}\,\{(z_0^2 - z_1^2)\frac{\partial}{\partial z_2} \wedge \frac{\partial}{\partial z_3} + z_0 z_3 \frac{\partial}{\partial z_0} \wedge \frac{\partial}{\partial z_2} - z_0 z_2 \frac{\partial}{\partial z_0} \wedge \frac{\partial}{\partial z_3} \\ & - z_1 z_3 \frac{\partial}{\partial z_1} \wedge \frac{\partial}{\partial z_2} + z_1 z_2 \frac{\partial}{\partial z_1} \wedge \frac{\partial}{\partial z_3}\}. \end{aligned}$$

Then $d\pi(w)$ is the Poisson structure on $\mathbb{CP}^3$. The 2-vector w is real since $f, g \in gl(2,\mathbb{H})\cap S^2\mathbb{C}^4$. Hence, the 2-vector w induces the Poisson structure on $\mathbb{HP}^1$.

Example 4.6. The polynomials

$$f = z_0^2 + z_1^2 + z_2^2 + z_3^2, \quad g = \sqrt{-1}\,(z_0 z_1 + z_2 z_3)$$

induce the 2-vector

$$\begin{aligned} w \;=\; & \sqrt{-1}\,\{(z_0^2 - z_1^2)\frac{\partial}{\partial z_2} \wedge \frac{\partial}{\partial z_3} + (z_2^2 - z_3^2)\frac{\partial}{\partial z_0} \wedge \frac{\partial}{\partial z_1} + (z_0 z_3 - z_1 z_2)\frac{\partial}{\partial z_0} \wedge \frac{\partial}{\partial z_2} \\ & + (z_1 z_3 - z_0 z_2)\frac{\partial}{\partial z_0} \wedge \frac{\partial}{\partial z_3} + (z_0 z_2 - z_1 z_3)\frac{\partial}{\partial z_1} \wedge \frac{\partial}{\partial z_2} + (z_1 z_2 - z_0 z_3)\frac{\partial}{\partial z_1} \wedge \frac{\partial}{\partial z_3}\} \end{aligned}$$

on $\mathbb{C}^4\backslash\{0\}$. Then w defines the Poisson structures on $\mathbb{CP}^3$ and $\mathbb{HP}^1$.

Similarly, there exist many global Poisson structures on $\mathbb{HP}^1$ of such type associated with holomorphic foliations on $\mathbb{CP}^3$.

Acknowledgements. The authors would like to thank Professor T. Taniguchi for his useful comments. They also would like to thank Professor R. Goto for his comment about the relationship between holomorphic foliations and holomorphic Poisson structures on $\mathbb{CP}^3$.

References

[1] Cerveau, D.; Lins Neto, A. Irreducible components of the space of holomorphic foliations of degree two in CP(n), $n \geq 3$, *Ann. of Math.* (2) **143** (1996), no. 3, 577–612

[2] Goto, R. Deformations of generalized complex and generalized Kähler structures, *J. Differential Geom.* **84** (2010), no. 3, 525–560.

[3] Hitchin, N. J. Instantons, Poisson structures and generalized Kähler geometry, *Comm. Math. Phys.* **265** (2006), no. 1, 131–164.

[4] Hitchin, N. J. Deformations of holomorphic Poisson manifolds, *Mosc. Math. J.* **12** (2012), no. 3, 567–591, 669.

Takayuki Moriyama
Department of Mathematics
Mie University
Mie 514-8507, Japan
E-mail: takayuki@edu.mie-u.ac.jp

Takashi Nitta
Department of Mathematics
Mie University
Mie 514-8507, Japan
E-mail: nitta@edu.mie-u.ac.jp